

Local monodromy of Drinfeld modules and t -motives

M. Mornev*

ETH Zürich

AMS Fall Eastern Virtual Sectional Meeting
2020

* Supported by ETH Zürich Postdoctoral Fellowship Program,
Marie Skłodowska-Curie Actions COFUND program

- K , a *local* field of *residual characteristic* p
- K^{sep} , a separable closure of K
- $G = \text{Gal}(K^{\text{sep}}/K)$

Structure of G

G absolute Galois group
 \cup
 I inertia
 \cup
 I^{0+} wild inertia

$$1 \rightarrow I \rightarrow G \rightarrow \text{Gal}(\bar{k}/k) \rightarrow 1$$

$$1 \rightarrow I^{0+} \rightarrow I \rightarrow \prod_{p' \neq p} \mathbb{Z}_{p'}(1) \rightarrow 1$$

$$I \twoheadrightarrow \mathbb{Z}/n\mathbb{Z}(1), \quad g \mapsto \frac{g(\sqrt[n]{\pi})}{\sqrt[n]{\pi}}, \quad (n, p) = 1, \quad \pi \text{ uniformizes } K$$

Local monodromy of ℓ -adic representations

$\ell \neq p$, a prime; \mathbb{Q}_ℓ -representations of G

The ℓ -adic monodromy theorem (Grothendieck)

Every ℓ -adic representation $\rho: G \rightarrow \mathrm{GL}(V)$ has the following properties:

- 1 The restriction of ρ to an open subgroup of I is unipotent.
- 2 The image $\rho(I^{0+})$ is finite.

In effect:

- $\rho(I^{0+}) = \{1\}$
- the group $I/I^{0+} = \prod_{p' \neq p} \mathbb{Z}_{p'}(1)$ acts through $\mathbb{Z}_\ell(1)$
- the action of $\mathbb{Z}_\ell(1) \cong \mathbb{G}_a(\mathbb{Z}_\ell)$ is *algebraic*

Weil-Deligne representations

$W \subset G$, the *Weil group*, preimage of $\mathbb{Z} \subset \widehat{\mathbb{Z}} = \text{Gal}(\bar{k}/k)$.

$\Phi \in W$ a Frobenius element; $t \in \mathbb{Z}_\ell(1)$ a generator $\rightsquigarrow \chi_t: I \rightarrow \mathbb{Z}_\ell$

WD: $V \mapsto (V, N)$

- $N: V(1) \rightarrow V$, $\rho(g) = \exp(\chi_t(g)N)$, $g \in I$ small
- $\text{WD}(\rho): \Phi^n g \mapsto \rho(\Phi^n g) \exp(-\chi_t(g)N)$

Theorem (Deligne)

The functor WD is an equivalence of categories of

- ℓ -adic representations of G ,
- Weil-Deligne representations in \mathbb{Q}_ℓ -vector spaces.

$X \rightarrow \text{Spec } K$ proper smooth

① *ℓ -independence conjecture:*

$\text{WD}(H^i(\bar{X}, \mathbb{Q}_\ell)) \otimes_{\mathbb{Q}_\ell} \mathbb{C}$ is independent of ℓ .

Known for abelian varieties and some other classes.

② *Weight-monodromy conjecture:*

The weight filtration on $H^i(\bar{X}, \mathbb{Q}_\ell)$ can be reconstructed from the monodromy operator.

F -representations

From now on: the characteristic of K is equal to p .

Instead of \mathbb{Q}_ℓ work with F , a local field of characteristic p .

E a Drinfeld module over $\text{Spec } K \rightsquigarrow V_p E$

Everything breaks down!

- 1 The image of I^{0+} is **typically** infinite.
- 2 The group I^{0+} is a free pro- p -group on countably many generators \rightsquigarrow An F -representation of I^{0+} is an essentially arbitrary infinite sequence of elements of $\text{GL}_n(F)$.

Game over? No.

Isocrystals

Fix a finite field \mathbb{F}_q of characteristic p ,
 \mathbb{F}_q -algebra structures on K and F .

$$\mathcal{E}_{K,F} = (\varinjlim_{n>0} K \otimes_{\mathbb{F}_q} \mathcal{O}_F/\mathfrak{m}_F^n) \otimes_{\mathcal{O}_F} F$$

$$F = \mathbb{F}_q((z)) \rightsquigarrow \mathcal{E}_{K,F} = K((z))$$

Endomorphism $\sigma: \mathcal{E}_{K,F} \rightarrow \mathcal{E}_{K,F}$ induced by the q -Frobenius of K .

Definition

An $\mathcal{E}_{K,F}$ -isocrystal is

- a finitely generated free $\mathcal{E}_{K,F}$ -module M
- equipped with an isomorphism $\tau_M^{\text{lin}}: \sigma^* M \xrightarrow{\sim} M$.

Morphisms are σ -equivariant morphisms of underlying modules.

Isocrystals and F -representations (1)

Tate module:

$$T(M) = (\mathcal{E}_{K^{\text{sep}}, F} \otimes_{\mathcal{E}_{K, F}} M)^T, \quad \tau: x \otimes m \mapsto \sigma(x) \otimes \tau_M^{\text{lin}}(1 \otimes m)$$

Definition

An isocrystal M is *unit-root* if $\dim_F T(M) = \text{rank}_{\mathcal{E}_{K, F}} M$.

Theorem (Katz)

The functor $M \mapsto T(M)$ is an equivalence of categories of

- unit-root $\mathcal{E}_{K, F}$ -isocrystals,
- F -representations of G .

Holds for every field K/\mathbb{F}_q , not just a local one.

Isocrystals and F -representations (2)

$$\mathcal{E}_{K,F}^+ = (\lim_{n>0} \mathcal{O}_K \otimes_{\mathbb{F}_q} \mathcal{O}_F/\mathfrak{m}_F^n) \otimes_{\mathcal{O}_F} F$$

$$F = \mathbb{F}_q((z)) \rightsquigarrow \mathcal{E}_{K,F}^+ = \mathcal{O}_K((z))$$

Theorem

The functor $M \mapsto T(\mathcal{E}_{K,F} \otimes_{\mathcal{E}_{K,F}^+} M)$ is an equivalence of cat-s of

- unit-root $\mathcal{E}_{K,F}^+$ -isocrystals,
- **unramified** F -representations of G .

Isocrystals and F -representations (3)

$$\mathcal{E}_{K,F}^b = K \otimes_{\mathcal{O}_K} \mathcal{E}_{K,F}^+$$

Theorem (M.)

The functor $M \mapsto \mathcal{E}_{K,F} \otimes_{\mathcal{E}_{K,F}^b} M$ is fully faithful on unit-root $\mathcal{E}_{K,F}^b$ -isocrystals.

Get a full subcategory of F -representations of G .

Proposition (M.)

For every

- Anderson A -module E over $\text{Spec } K$,
 - prime $\mathfrak{p} \subset A$ different from the residual characteristic of E
- the Tate module $V_{\mathfrak{p}}E$ arises from $\mathcal{E}_{K,F}^b$.

$$\mathfrak{p} \neq \text{res. char.}(E) \sim \ell \neq p$$

Upper index ramification filtration I^v , $v \in \mathbb{Q}_{\geq 0}$
 $I^{0+} = \text{closure of } \bigcup_{v>0} I^v$

Conjecture (the z -adic monodromy theorem)

An F -representation $\rho: G \rightarrow \text{GL}(V)$ arises from $\mathcal{E}_{K,F}^b$ if and only if

- 1 the restriction of ρ to an open subgroup of I is unipotent,
- 2 $\rho(I^v) = \{1\}$ for $v \gg 0$.

In the ℓ -adic case: $\rho(I^{0+})$ is finite.

Property (1) holds for Anderson modules by Gardeyn's analytic monodromy theorem.

Theorem (M.)

Property (2) holds for Drinfeld modules.

Under development:

- analog of Weil-Deligne construction,
- ℓ -independence conjecture,
- weight-monodromy conjecture.

$\rho: G \rightarrow \mathrm{GL}(V)$, an F -representation of G

$V_0 \subset V$ an unramified representation such that V/V_0 is also unramified. This holds for Drinfeld modules.

p -adic monodromy theorem: $\rho(I^v) = \{1\}$ for some $v \gg 0$.

- J^v , an abelian quotient of I depending only on v ,
- J^v is finitely generated over $\mathbb{F}_p[[\phi]]$ where $1 + \phi$ acts as the Frobenius
- Finite-dimensional $\mathbb{F}_p((\phi))$ -vector space
 $H = \mathrm{Hom}_F(V_0, V/V_0)$.
- $\rho|_I$ is encoded by an $\mathbb{F}_p[[\phi]]$ -linear homomorphism

$$\chi: J^v \rightarrow H$$